

## **MOXI™ LASER AFTER CARE (PAGE 1 of 2)**

Redness is normal and expected. Redness generally increases in intensity the first few days after treatment with day 3 being most intense. Redness can persist for up to 7 days depending on the aggressiveness of the treatment.

Pinpoint bleeding may occur. This can last for a few hours - 12 hours depending on the treatment depth and may be isolated to certain treated areas.

Immediately after treatment, swelling is common and expected. Use of a cold compress or ice packs will help to relieve the swelling. To avoid further swelling, you may choose to sleep in an upright position the first night after the treatment. The first morning post treatment is when swelling is more prevalent, especially under the eyes. Swelling may last 2-4 days depending on the aggressiveness of the treatment.

The treated area may be extremely warm for 12-24 hours after the treatment. Cold compresses or ice packs may provide comfort during this time.

On the 2<sup>nd</sup> or 3<sup>rd</sup> day after treatment, you may increasingly notice tiny dark spots and bronzed appearance to the treated skin. This is called the MENDS (microscopic epidermal necrotic debris). In individuals with heavily pigmented skin, or in areas where sun damage has produced pigmented lesions, the microscopic wounds, known as MENDS, contain large amounts of melanin. Because there are so many MENDS, they can make the skin look bronzed and small areas appear crusted (do not try to scratch them off). This is part of the healing process where treated tissue is working its way out of your body as new fresh skin is regenerated. During this time, your skin will be very dry and have a sandpaper texture and will begin to flake and peel. If the face is treated, it could take 5-7 days for the peeling process to be completed. If the neck, chest or anywhere on the body was treated, this process could take up to 2 weeks.

After the peeling process is complete, your skin will have a rosy, pink glow that will gradually resolve. You can apply makeup after 48 hours of the procedure.

Antiviral medication: If an antiviral was prescribed for you, continue to take as directed. Most commonly, it will be Valacyclovir 500-1000 mg 2 times a day for a total of 7 days, preferably starting 3 days prior to the procedure.

Medication for swelling: If you were instructed to do so, use Claritin or Zyrtec once a day starting the week prior to the treatment and continuing for 3 days after the treatment along with Benadryl 25 mg at night starting the night of the treatment for 2-3 nights.

Post treatment discomfort may be relieved by over the counter oral pain relievers, i.e. Extra Strength Tylenol or Ibuprofen if recommended by your practitioner.

Itching may be experienced during the healing phase and is completely normal. Oral Benadryl may help itching but can cause drowsiness. DO NOT scratch the treated area as scarring and pigmentation complications can occur.

**MOXI™ LASER AFTER CARE (PAGE 2 of 2)**

**Keep your skin moist with Cicalfate**, Aquaphor or other moisturizing cream that your practitioner recommends. It should be reapplied as needed whenever your skin feels dry. The use of a moisturizer with petroleum may be suited for individuals who receive more graduated coverage while the cream may be sufficient on less aggressive treatments.

Cleanse the skin two times a day with plain, lukewarm water and a gentle cleanser, e.g. Cetaphil, beginning the morning after the treatment. Use your hands to gently apply the cleanser and water and finish by patting dry with a soft cloth. **DO NOT** rub, scrub, use an exfoliant or a skin care brush e.g. Clarisonic in the treated area. Doing so could result in scarring and pigmentation complications.

Peeling and flaking generally occur within 24 hours post treatment and should be allowed to come off naturally. **DO NOT PICK, RUB, OR FORCE OFF ANY SKIN DURING THE HEALING PROCESS, THIS COULD RESULT IN SCARRING, PIGMENTATION COMPLICATIONS AND INFECTION!** Gently washing the skin more frequently will help to promote the peeling process.

**Sunscreen is a MUST** and should be used daily beginning the day of treatment and used vigilantly for up to 3 months post procedure. Use recommended sunscreen with broadband protection (UVA and UVB) and a sun protection factor (SPF) of 30. Apply sunscreen 20 minutes before sun exposure. Reapply sunscreen every 2 hours. If direct sun exposure is necessary, wear a hat and clothing that covers the treated area. If treated area is exposed to sun (direct or indirect) blistering, scarring, hyperpigmentation, or hypo-pigmentation can / will occur. Sun exposure should be avoided for 2 weeks post treatment.

Typically, **after** the peeling process is complete, make up can be worn.

When showering in the next few days, be sure to avoid getting shampoo directly on the treated area. You may consider taking your shower with the back of your head to the water to avoid directly hitting your face with the full force stream of the water.

Avoid strenuous exercise and sweating until after skin has healed.

There may be some degree of swelling immediately post treatment; however, if you have excessive swelling or any of the following signs of infection, you should contact the office immediately including drainage that looks like pus; increased warmth at or around the treated area; fever of 101.5 or greater; extreme itching.

Usually 3-6 Moxi treatments are required to achieve desired goals. Your provider will help guide you.

Make a follow up appointment for 2 weeks after your treatment with your provider.

Your final results will not be realized for up to 3 months post final procedure. Be patient.

Call RejuvenationMD with any questions.