

BBL ACNE TREATMENT AFTER CARE

Patient response can vary after a BBL Acne treatment. Erythema (redness) is usually noted within a few minutes after the completion of the procedure. A slight sun burn sensation in and around the area treated is also normal and expected. These reactions tend to subside within 1-4 hours after the treatment. The treated area must be treated with care. **BE GENTLE!** Do not scratch or pick at your skin.

Until sensitivity has completely subsided, avoid all the following:

- Use of scented lotions or soaps, exfoliant creams (Retin-A, glycolic/salicylic and alpha-hydroxy acids), acne creams or gels, loofa sponges and aggressive scrubbing
- Hot or cold water - wash with tepid water
- Shaving
- Swimming pools and spas with multiple chemicals/chlorine
- Activities that cause excessive perspiration

A cold compress or an ice pack can be used to provide comfort if the treated area is especially warm. This is rarely needed but if so, usually within the first 4 hours after the treatment.

Makeup can be applied immediately after treatment as long as skin is intact and there are no signs of blistering.

Vitamin E or Aloe Vera applied to the treatment area may provide a soothing effect.

Post treatment discomfort is typically minimal but if the area is very uncomfortable, oral pain relievers, i.e., Extra Strength Tylenol or Advil, may be used.

If the skin is broken or a blister appears, apply an antibiotic ointment, and **contact** the office immediately.

Strictly avoid any sun exposure to the treated area for a minimum of 7 days after the procedure. The treated area is more prone to sunburn and pigmentation change. Keep the area covered and use a sun block with a protection factor of 30+ reapplying the sun block every two hours. Delayed blistering secondary to sun exposure has been noted up to 72 hours post treatment.