

EZ PRF/EZ GEL INJECTIONS PRE- AND POST CARE

Pre-Treatment Instructions:

- Avoid the following for at least 1 week prior to your procedure:
 - o NSAIDS (Ibuprofen, Motrin, Aleve, Naprosyn, Aspirin)—Tylenol is allowed.
 - Certain nutritional supplements (Arnica, Gingko, Garlic, Vitamin E, Vitamin A, Flax Oil,
 Curcumin, and other anti-inflammatory nutrients)
 - Systemic Steroids (prednisone, dexamethasone)
 - Alcohol and Cigarettes
- Hydrate well the day before and the day of the procedure for ease of blood draw.

On the Treatment Day:

- Blood is drawn and PRF is processed.
- Topical anesthetic is applied to the treatment area for 20-30 minutes if needed.
- Filler may be injected into the areas of concern first which can be followed by the PRP/PRF/EZ
 Gel.
- If we are only using PRP/PRF/EZ Gel, it will be injected into the areas of concern at this time.
- Fillers are injected when a defect is being augmented (indentations, inverted nipple).
- PRP and PPP (Platelet Poor Plasma) is then injected in multiple areas of the breasts to facilitate rejuvenation and regeneration of tissue.
- There may be redness, swelling and bruising after the procedure.

Post-Treatment Instructions:

- Expect mild inflammation, redness, swelling for a few days.
- Bruising may last several weeks. You may apply arnica gel or cream to the area to help. Do <u>not</u> take Arnica internally, however.
- If there is significant swelling, you can use oral antihistamines such as Claritin/Allegra/Zyrtec (or generic equivalents) during the day and/or Benadryl at night.
- Avoid manipulating or pressing on the treated area for 48 hours (including having other facial treatments done). If you are concerned with any lumps felt, please call the office for a follow up appointment within a week.
- Try to sleep on an incline and not on your belly.
- Avoid any massages where you are pushing your face into a massage table.
- For at least 1 week, avoid NSAIDS (Ibuprofen, Motrin, Aleve, Naprosyn, Aspirin), certain nutritional supplements (Arnica, Gingko, Garlic, Vitamin E, Vitamin A, Flax Oil, Curcumin and other anti-inflammatory nutrients), systemic steroids (prednisone, dexamethasone), alcohol and cigarettes.
- Eat healthy and hydrate very well (at least 64 ounces of water daily).